

Available online at: <https://ijact.in>

Date of Submission	30/05/2019
Date of Acceptance	19/06/2019
Date of Publication	03/07/2019
Page numbers	3197-3200 (4 Pages)

Cite This Paper: Natalja O. Gordeeva, Ekaterina N. Manaeva, Ilona M. Primak, Irina I. Palasheva.(2019). Use of finite difference method for numerical solution of three-dimensional heat transfer fractional differential equation, 8(6), COMPUSOFT, An International Journal of Advanced Computer Technology. PP. 3197-3200.

This work is licensed under Creative Commons Attribution 4.0 International License.



ISSN:2320-0790

USE OF FINITE DIFFERENCE METHOD FOR NUMERICAL SOLUTION OF THREE-DIMENSIONAL HEAT TRANSFER FRACTIONAL DIFFERENTIAL EQUATION

Natalja O. Gordeeva*, Ekaterina N. Manaeva, Ilona M. Primak, Irina I. Palasheva

Belgorod State National Research University, Starooskolsky Branch 18 Solnechny microdistrict, 4
Stary Oskol, 309502, Russia

Abstract: The paper proposes a numerical solution for the mixed problem concerning a three-dimensional heat transfer fractional differential equation, based on the finite difference method. To solve this problem, an explicit difference scheme described in the paper is used. The stability of a proposed difference scheme is proved. The case of homogeneous medium and a square grid is considered.

Keywords: fractional differential equations; heat and mass transfer equation; numerical methods; approximation with fractional derivatives.

I. INTRODUCTION

The development of the fractional integration and differentiation apparatus technique is of interest both in terms of the development of the fractional integrodifferentiation theory itself and for solving practical problems.

Fractional integrals and derivatives and fractional integrodifferential equations find their application in theoretical physics, mechanics, applied mathematics, where their application allows a deeper understanding of known results and obtaining a new class of solutions, and cover a wide range of problems not previously explained from the standpoint of traditional approaches. Fractional integrodifferentiation is a powerful tool for describing

physical systems with memory and non locality characteristics.

Despite the long history of the development of the fractional differentiation mathematical apparatus technique, analytical methods for solving fractional derivative equations have been obtained only for a narrow circle of problems [1–5]. At the same time, numerical methods for solving such problems are being actively developed now [6–8].

II. METHOD

Formulation of the problem

Parabolic equations with fractional derivatives [9-11] are used in the study of heat transfer problems. We consider a

non stationary three-dimensional heat transfer equation with fractional derivatives.

$$D_t^\alpha T(t, x, y, z) = C_x D_x^\beta T(t, x, y, z) + C_y D_y^\beta T(t, x, y, z) + C_z D_z^\beta T(t, x, y, z),$$

Where C_x, C_y, C_z are heat conductivity coefficients, $0 < \alpha \leq 1, 1 < \beta \leq 2$.

We find a solution to this equation $T(t, x, y, z) \in C^3(D)$ ($D = \{(x, y, z, t) : 0 \leq x \leq a, 0 \leq y \leq b, 0 \leq z \leq c, 0 \leq t \leq T\}$) satisfying the initial and boundary conditions:

$$T(0, x, y, z) = \psi(x, y, z), x, y, z \in G \quad (2)$$

$$T(t, x, y, z)|_\Gamma = \varphi(x, y, z), x, y, z \in G \quad (3)$$

Where G is a parallelepiped with the border Γ . The equation contains partial fractional derivatives of Riemann-Liouville:

$$D_x^\beta T(t, x, y, z) = \left(\frac{\partial}{\partial x}\right)^2 I^{2-\beta} T(t, x, y, z)$$

and the partial fractional integral of Riemann-Liouville with respect to the corresponding variables:

$$I_x^\beta T(t, x, y, z) = \frac{1}{\Gamma(\beta)} \int_0^x T(t, s, y, z) (x-s)^{1-\beta} ds.$$

III. RESULT AND DISCUSSION

We describe the numerical scheme used to solve the equation and introduce a grid in the region D :

We divide the region $[0; a]$ to $L+1$ segments with nodes $x_l = l\Delta x, (l = 0, \dots, L)$ where $\Delta x = \frac{a}{L}$ is a step along x .

Then we divide the region $[0; b]$ to $M+1$ segments with nodes $y_m = m\Delta y, (m = 0, \dots, M)$ where $\Delta y = \frac{b}{M}$ is a step along y . Then we divide the region $[0; c]$ to $N+1$ segments with nodes

$z_n = n\Delta z, (n = 0, \dots, N)$ where $\Delta z = \frac{c}{N}$ is a step along

z . Split the time interval $[0; T]$ into parts with a step Δt and time nodes $t_k = k\Delta t, (k = 0, \dots, K)$.

For convenience, we denote $T_{k,l,m,n} = T(t_k, x_l, y_m, z_n)$ and use the Grünwald formula [12] to approximate the fractional Riemann – Liouville derivative.

It is known [12, p. 280] that for any function $f(t)$ which allows decomposition in power series, a fractional derivative of order $1-\gamma$ at any point of convergence of this series can be written as a Grünwald-Letnikov derivative:

$$D_t^{1-\gamma} f(t) = \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t^{1-\gamma}} \sum_{k=0}^{\lfloor \frac{t}{\Delta t} \rfloor} \omega_k^{(1-\gamma)} f(t - k\Delta t) \quad (4)$$

Where $\omega_0 = 1, \omega_k = \frac{(-1)^k \beta(\beta-1)\dots(\beta-k+1)}{k!}$

If the function is continuous and its derivative is integrable over the segment $[0; t]$, then Riemann-Liouville and Grünwald-Letnikov derivatives of any order $0 < 1-\gamma < 1$ exist and match for any point in time from $[0; t]$ [9].

The Grünwald-Letnikov definition allows us to numerically find the Riemann-Liouville derivative:

$$D_t^{1-\gamma} f(t) \approx \frac{1}{\Delta t^{1-\gamma}} \sum_{k=0}^{\lfloor \frac{t}{\Delta t} \rfloor} \omega_k^{(1-\gamma)} f(t - k\Delta t) \quad (5)$$

Using the formula (5) for the fractional Riemann-Liouville derivatives with respect to the spatial variables, we obtain:

$$D_x^\beta T(t, x, y, z)|_L = \frac{1}{\Delta x^\beta} \sum_{j=0}^{L+1} \omega_j^\beta T(t, x - j\Delta x, y, z)$$

$$D_y^\beta T(t, x, y, z)|_M = \frac{1}{\Delta y^\beta} \sum_{j=0}^{M+1} \omega_j^\beta T(t, x, y - j\Delta y, z),$$

$$D_z^\beta T(t, x, y, z)|_N = \frac{1}{\Delta z^\beta} \sum_{j=0}^{N+1} \omega_j^\beta T(t, x, y, z - j\Delta z).$$

Considering that $x_{L-j} = x_L - j\Delta x, y_{M-j} = y_M - j\Delta y, z_{N-j} = z_N - j\Delta z$, we will present Riemann-Liouville derivative at $0 < \alpha \leq 1$ in the form [6]:

$$D_0^\alpha T(t, x, y, z)|_k = \frac{1}{\Gamma(1-\alpha)} \left(\frac{T(t_k, x, y, z)}{(t_{k+1} - t_k)^\alpha} + \int_{t_k}^{t_{k+1}} \frac{T'(s, x, y, z) ds}{(t_{k+1} - s)^\alpha} \right)$$

(6)

Where $\Gamma(\alpha) = \int_0^\infty x^{\alpha-1} e^{-x} dx$

We present the derivative $\frac{dT(s, x, y, z)}{ds}$ on the segment $[t_k; t_{k+1}]$ in the form of a finite difference [13, p.47]:

$$\left(\frac{dT}{ds}\right)_k \approx \frac{T(t_{k+1}, x, y, z) - T(t_k, x, y, z)}{\Delta t} \tag{7}$$

In view of (6), the Riemann-Liouville derivative of fractional order $[t_k; t_{k+1}]$ at the interval can be approximated by the finite difference:

$$D_0^\alpha T(t, x, y, z)|_m = \frac{1}{\Gamma(1-\alpha)} \left(\frac{T(t_m, x, y, z) - T(t_{m-1}, x, y, z)}{\Delta t^\alpha} + \frac{T(t_{m+1}, x, y, z) - T(t_m, x, y, z)}{\Delta t} \int_m^{m+1} \frac{ds}{(t_{m+1} - s)^\alpha} \right) = \frac{T(t_{m+1}, x, y, z) - \alpha T(t_m, x, y, z)}{\Gamma(1-\alpha)(1-\alpha)\Delta t^\alpha}$$

As a result, for the heat transfer equation (1), we write the explicit difference scheme:

$$\frac{T_{k+1,j,m,n} - \alpha T_{k,j,m,n}}{\Gamma(1-\alpha)(1-\alpha)\Delta t^\alpha} = \frac{C_x}{\Delta x^\beta} \sum_{j=0}^{i+1} \omega_j^\beta T_{k,j-1,m,n} + \frac{C_y}{\Delta y^\beta} \sum_{j=0}^{m+1} \omega_j^\beta T_{k,j,m-j+1,n} + \frac{C_z}{\Delta z^\beta} \sum_{j=0}^{n+1} \omega_j^\beta T_{k,j,m,n-j+1}$$

or taking into account the values ω_j

$$\frac{T_{k+1,j,m,n} - \alpha T_{k,j,m,n}}{\Gamma(1-\alpha)(1-\alpha)\Delta t^\alpha} = \frac{C_x}{\Delta x^\beta} (T_{k,j+1,m,n} - \beta T_{k,j,m,n} + \sum_{j=2}^{j+1} \omega_j^\beta T_{k,j-j+1,m,n}) + \frac{C_y}{\Delta y^\beta} (T_{k,j,m+1,n} - \beta T_{k,j,m,n} + \sum_{j=2}^{m+1} \omega_j^\beta T_{k,j,m-j+1,n}) + \frac{C_z}{\Delta z^\beta} (T_{k,j,m,n+1} - \beta T_{k,j,m,n} + \sum_{j=2}^{n+1} \omega_j^\beta T_{k,j,m,n-j+1})$$

The decision $T(t_{k+1}, x, y, z)$ can be represented as:

$$T(t_{k+1}, x, y, z) = ST(t_k, x, y, z),$$

Where S is the operator for a transition from onetime layer to another.

Then we find out the stability of the scheme. We have

$$\|T(t_{k+1}, x, y, z)\| = \|ST(t_k, x, y, z)\| \leq \|S\| \cdot \|T(t_k, x, y, z)\| \text{ and } \|T(t_{k+1}, x, y, z)\| \leq \|T(t_k, x, y, z)\| \leq \dots \leq \|T(t_0, x, y, z)\|$$

From where it follows, that the initial disturbances fade out. Given the initial data, the stability condition of the difference scheme (8) is [14]. This means that the spectrum of the operator lies inside a circle of a unit radius on the complex plane. From there, where λ is any eigenvalue of the transition operator. We find the eigenvalues of the transition operator, for this, we will

present the solution $T(t_k, x, y, z)$ in the form of disturbance.

$$T(t_k, x, y, z) = \lambda^k \exp(i\omega_x \Delta x) \exp(i\omega_y \Delta y) \exp(i\omega_z \Delta z), \tag{9}$$

Where i is an imaginary unit, and λ is an eigenvalue of the transition operator [14].

We substitute equality (9) into (8), and then obtain the estimate for the eigenvalues of the transition operator:

$$\lambda \leq \alpha - \frac{\Gamma(2-\alpha)C_x \Delta t^\alpha}{\Delta x^\beta} (4 \sin^2 \frac{\omega_x \Delta x}{2} + \beta - 2) - \frac{\Gamma(2-\alpha)C_y \Delta t^\alpha}{\Delta y^\beta} (4 \sin^2 \frac{\omega_y \Delta y}{2} + \beta - 2) - \frac{\Gamma(2-\alpha)C_z \Delta t^\alpha}{\Delta z^\beta} (4 \sin^2 \frac{\omega_z \Delta z}{2} + \beta - 2). \tag{10}$$

From where we have, that all the eigenvalues of the transition operator do not exceed 1 in absolute value. We get:

$$0 \leq \frac{\Gamma(2-\alpha)C_x \Delta t^\alpha}{\Delta x^\beta} + \frac{\Gamma(2-\alpha)C_y \Delta t^\alpha}{\Delta y^\beta} + \frac{\Gamma(2-\alpha)C_z \Delta t^\alpha}{\Delta z^\beta} \leq \frac{\alpha + 1}{\beta + 2}.$$

IV. CONCLUSION

Thus, the following assertion is proved. Statement 1.

Explicit difference scheme

$$\frac{T_{k+1,j,m,n} - \alpha T_{k,j,m,n}}{\Gamma(1-\alpha)(1-\alpha)\Delta t^\alpha} = \frac{C_x}{\Delta x^\beta} \sum_{j=0}^{i+1} \omega_j^\beta T_{k,j-1,m,n} + \frac{C_y}{\Delta y^\beta} \sum_{j=0}^{m+1} \omega_j^\beta T_{k,j,m-j+1,n} + \frac{C_z}{\Delta z^\beta} \sum_{j=0}^{n+1} \omega_j^\beta T_{k,j,m,n-j+1}$$

Is stable if

$$\Delta t^\alpha \left(\frac{C_x}{\Delta x^\beta} + \frac{C_y}{\Delta y^\beta} + \frac{C_z}{\Delta z^\beta} \right) \leq \frac{\alpha + 1}{(\beta + 2)\Gamma(2-\alpha)}.$$

We consider the case of a homogeneous medium $C_x = C_y = C_z = C$ and a square grid when

$\Delta x = \Delta y = \Delta z$. The difference scheme (8) will be:

$$\frac{T_{k+1,j,m,n} - \alpha T_{k,j,m,n}}{\Gamma(1-\alpha)(1-\alpha)\Delta t^\alpha} = \frac{C}{\Delta x^\beta} \left((T_{k,j+1,m,n} + \sum_{j=2}^{j+1} \omega_j^\beta T_{k,j-j+1,m,n}) + (T_{k,j,m+1,n} + \sum_{j=2}^{m+1} \omega_j^\beta T_{k,j,m-j+1,n}) + (T_{k,j,m,n+1} + \sum_{j=2}^{n+1} \omega_j^\beta T_{k,j,m,n-j+1}) - 3\beta T_{k,j,m,n} \right) \tag{12}$$

An explicit difference scheme (12) is stable when:

$$\Delta t^\alpha \leq \frac{\Delta x^\beta (\alpha + 1)}{3C(\beta + 2)\Gamma(2-\alpha)}.$$

A great difficulty is inherent in a computational experiment based on the obtained difference scheme due to an increase in the volume of computations as the value of the time variable increases. As a result, the computation time of even a one-dimensional problem on a relatively coarse grid takes several hours. Therefore, for such calculations, it seems reasonable to use multiprocessor computing, which entails the need to appropriately adapt existing methods and algorithms.

V. REFERENCES

- [1] Golovizin V.M., Kisilev V.P., Korotkin I.A. Numerical methods for solving the fractional diffusion equation in the one-dimensional case: Preprint No. IBRAE-2002-10. Moscow: IBRAE RAS, 2002.
- [2] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo. 2006. Theory and Applications of Fractional Differential Equations, Volume 204 (North-Holland Mathematics Studies). Elsevier Science Inc., New York, NY, USA. ISBN:0444518320.
- [3] Bajlekova E. Fractional Evolution Equations in Banach Spaces. Ph. D. Thesis. Eindhoven University of Technology. 2001. — 117 p. url: <https://pure.tue.nl/ws/files/2442305/200113270.pdf> (Accessed on 30.12.2018)
- [4] Glushak A.V., Primak I.M. Boundary value problems for abstract differential equations with fractional derivatives / Scientific Bulletin of the BelSU. Series: Mathematics. Physics. ISSN: 2075-4639, 2011. - №17 (112). - Issue 24. - Pp.125-140.
- [5] Primak I.M. Boundary value problems for abstract fractional differential equations with a bounded operator./Scientific Bulletin of the BelSU. Series: Mathematics. Physics. ISSN: 2075-4639, 2013. - №5 (148). - Issue 30. - Pp.98-106.
- [6] Nakhushiev A.M. Fractional calculus and its application. Moscow: Fizmatlit, 2003. 272 p.
- [7] Beibalaev V.D., Shabanova M.R. A numerical method for solving a boundary value problem for a two-dimensional heat transfer equation with fractional derivatives. Herald of the Samara State Technical University. Series: Physics and mathematics. - 2010. - № 5 (21). - pp. 244–251.
- [8] Accessed from: http://www.mathnet.ru/php/archive.phtml?wshow=paper&jrnid=vsgtu&paperid=776&option_lang=eng (Accessed on 14.01.2019)
- [9] Taukenova F. I., Shkhanukov-Lafishev M.Kh. Difference methods for solving boundary value problems for fractional-order differential equations // Comput. Math. Math. Phys., 2006. — Vol. 46, No. 10. — P. 1785–1795. url: <https://link.springer.com/article/10.1134/S0965542506100149#citeas> (accessed on 22.02.2019)
- [10] Lynch V.E., Carreras B.A., del-Castill-Negrete D., Ferreira-Mejias K.M., Hicks H.R. Numerical methods for the solution of partial differential equations of fractional order // J. Comput. Phys., 2003. — Vol. 192, No. 2. — Pp. 406–421
- [11] Babenko Yu.I. The fractional differentiation method in applied problems of heat and mass transfer theory. - SPb: NPO Professional, 2009, 584 p.
- [12] S. B. Yuste, and L. Acedo. “An Explicit Finite Difference Method and a New Von Neumann-Type Stability Analysis for Fractional Diffusion Equations.” SIAM Journal on Numerical Analysis, vol. 42, no. 5, 2005, pp. 1862–1874. JSTOR, www.jstor.org/stable/4101366
- [13] Samko S.G., Kilbas A.A., Marichev O.I. Fractional integrals and derivatives and some of their applications. Minsk: Science and technology, 1987, 688 p.
- [14] Samko S.G., Kilbas, A.A. and O.I. Marichev: Fractional Integrals and Derivatives,; Theory and Applications, Gordon and Breach, Amsterdam 1993. [Engl. Transl. from Russian, Integrals and Derivatives of Fractional Order and Some of Their Applications, Nauka i Tekhnika, Minsk 1987]”
- [15] Daletsky Yu.L., Krein M.G. Stability of solutions of differential equations in a Banach space. Nauka, Moscow, 1970. - 536 p.